

12-6-8



200132

NURSERY POLYON

Average Media Temperatures	60°F	70°F	80°F	90°F
Longevity in Months	3+	2-3	1-2	

Application Rates

Incorporation

Uniformly mix (incorporate) Custom Nursery Fertilizer into potting media at following rates using trials and grower experience to obtain optimum performance. The addition of dolomitic lime-stone, additional micronutrients or other media amendments may be needed.

POUNDS PER CUBIC YARD			
Low	Medium	High	South FL
8.8	10.8	13.1	0

Topdressing - Rates Per Container

Uniformly apply product away from the base of plant.

(gallons)	Low	Medium	High	South FL
Trade G (g)	5	7	8	0
1 (g)	8	10	12	0
2 (g)	19	23	29	0
3 (g)	34	42	51	0
5 (g)	46	56	69	0
7 (g)	69	84	102	0
10 (g)	86	105	128	0
15 (g)	96	117	143	0
20 (g)	127	155	189	0
25 sqt. (g)	188	230	280	0
25 (g)	188	230	280	0
#35 (g)	210	256	312	0

**NET WT.
50 LBS.
(22.7 KG)**

MANUFACTURED FOR HARRELL'S LLC. (F352) 720 Kraft Road, Lakeland, FL 33801 - (863) 687-2774 - (800) 282-8007

Distributed by: Harrell's, LLC. P.O. Box 187, Sylacauga, AL 35151 - (256) 245-5559 - (800) 282-8007

The GREEN color of the Polyon® capsule and RLC (Reactive Layers Coating) are trademarks of RLC Technologies, L.L.C. POLYON® and POLYON NPK® are registered trademarks of RLC Technologies.

GUARANTEED ANALYSIS

* Total Nitrogen (N)	12.0000%
3.4700% Ammoniacal Nitrogen	
0.0900% Water-Soluble Nitrogen	
7.6100% Urea Nitrogen	
0.8300% Water Insoluble Nitrogen	
Available Phosphate (P2O5)	6.0000%
Soluble Potash (K2O)	8.0000%
Magnesium (Mg)	3.6000%
2.244% Water Soluble Magnesium (Mg)	
Copper (Cu)	0.1250%
0.001% Water Soluble Copper (Cu)	
Iron (Fe)	1.3560%
0.064% Water Soluble Iron (Fe)	
0.125% Iron (Chelated)	
Manganese (Mn)	0.3750%
0.009% Water Soluble Manganese (Mn)	
Zinc (Zn)	0.1250%
0.001% Water Soluble Zinc (Zn)	
Derived From: Activated Sludge, Ammoniated Phosphate, Polymer Coated Urea, Sulfate of Ammonia, Sulfate of Potash, Sulfate of Potash-Magnesia, Copper Sulfate, Copper Oxide, Iron EDTA, Iron Oxide, Iron Sulfate, Magnesium Oxide, Magnesium Sulfate, Manganese Oxide, Manganese Sulfate, Zinc Sulfate, Zinc Oxide	
* 6.8% slow release NITROGEN derived from Activated Sludge, Polymer Coated Urea	
Chlorine, Not more than.....	.0700%

Density - 64 lb./cu. ft.)

Directions for Use

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to, a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

CONVERSION TABLE

Rounded Measure	Gram:
1 teaspoon (tsp)	7.1
1 tablespoon (tbs.)	22.0
1/4 cup	61.4
1/2 cup	141.5

†The gram weights are approximate. Actual weight in your measuring should be determined for the fertilizer being applied.

Harrell's SPOON SIZE/RATE

Spoon Size	Grams
1	10
2	15
3	20
4	26
5	38
6	54
7	70
8	88

KEEP OUT OF REACH
OF CHILDREN
CAUTION
HARMFUL IF
SWALLOWED
AVOID INHALATION
OF PRODUCT

APPLICATION PRECAUTIONS

- A trial of this product on a limited scale application methods, and desired rates is recommended. Avoid application to plants under stress.
- Do not incorporate into media prior to steam sterilization.
- For mixes containing soil, reduce application rate by approximately 25%.
- When using supplemental liquid feed, reduce the rate of this formulation accordingly.
- If mixed media is not used within two weeks, leach thoroughly before using.
- To avoid buildup of soluble salts, occasional leaching may be necessary.

1001-0784 051002